

# Leukaemia Section

## Short Communication

### t(11;16)(q23;p13)

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France

Published in Atlas Database: February 1998

Online updated version: <http://AtlasGeneticsOncology.org/Anomalies/t1116q23p13ID1120.html>

DOI: 10.4267/2042/37419

This work is licensed under a Creative Commons Attribution-Non-commercial-No Derivative Works 2.0 France Licence.  
© 1998 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

## Clinics and pathology

### Disease

ANLL/MDS: only treatment related leukaemias cases so far (in other 11q23 translocations, most cases occur in de novo acute leukaemia).

### Phenotype / cell stem origin

M4, M2 ANLL; CMML and RAEBT, although MDS is otherwise rarely seen in 11q23 translocations; the fusion gene is found in all the mature monocytes, in some of the granulocytes and erythroblasts, not in the lymphocytes.

### Epidemiology

13 available cases; most cases are children cases; median age is 10-14 yrs, range is 2-74 yrs; sex ratio is balanced.

### Clinics

Secondary to antitopoisomerase II drugs (etoposide or teniposide, but also doxorubicin); this secondary malignancy occurs within 6-60 mths (median 20 mths); the primary malignancy was a t(8;21)(q22;q22)/M2-ANLL in 2 cases.

### Prognosis

Yet unknown.

## Cytogenetics

### Additional anomalies

Are found in 8 of 11 cases; variable, except the unexpected recurrence of 1p36.1 involvement.

## Genes involved and Proteins

### MLL

Location: 11q23

### DNA / RNA

21 exons, spanning over 100 kb; 13-15 kb mRNA.

### Protein

431 kDa; contains two DNA binding motifs (a AT hook, and Zinc fingers), a DNA methyl transferase motif, a bromodomain; transcriptional regulatory factor; nuclear localisation.

### CBP

Location: 16p13

### Protein

Nuclear localisation; transcriptional adaptor/coactivator: binds CREB; has histone acetyltransferase activity.

## Results of the chromosomal anomaly

### Hybrid gene

### Description

5' MLL - 3' CBP

### Fusion protein

### Description

N-term AT hook and DNA methyltransferase from MLL fused to most of CBP starting with the bromodomain of CBP -or even more in N-term with the CREB binding domain- and also comprising the cystein/histidine rich and the glutamine rich domains of CBP in C-term around 1400 amino acids from MLL; the reciprocal CBP-MLL may or may not be expressed.

### Oncogenesis

May promote histone acetylation of genomic regions targeted by the MLL AT-hooks; may loose CBP cell cycle inhibition capability.

## References

Satake N, Ishida Y, Otoh Y, Hinohara S, Kobayashi H, Sakashita A, Maseki N, Kaneko Y. Novel MLL-CBP fusion transcript in therapy-related chronic myelomonocytic leukemia with a t(11;16)(q23;p13) chromosome translocation. *Genes Chromosomes Cancer* 1997 Sep;20(1):60-3.

Taki T, Sako M, Tsuchida M, Hayashi Y. The t(11;16)(q23;p13) The t(11;16)(q23;p13) translocation in myelodysplastic syndrome fuses the MLL gene to the CBP gene.. *Blood* 1997 Jun 1;89(11):3945-50.

Rowley JD, Reshmi S, Sobulo O, Musvee T, Anastasi J, Raimondi S, Schneider NR, Barredo JC, Cantu ES, Schlegelberger B, Behm F, Doggett NA, Borrow J, Zeleznik-Le N. All patients with the T(11;16)(q23;p13.3) that involves MLL

and CBP have treatment-related hematologic disorders. *Blood* 1997 Jul 15;90(2):535-41.

Sobulo OM, Borrow J, Tomek R, Reshmi S, Harden A, Schlegelberger B, Housman D, Doggett NA, Rowley JD, Zeleznik-Le NJ. MLL is fused to CBP, a histone acetyltransferase, in therapy-related acute myeloid leukemia with a t(11;16)(q23;p13.3). *Proc Natl Acad Sci USA* 1997 Aug 5;94(16):8732-7.

---

*This article should be referenced as such:*

Huret JL. t(11;16)(q23;p13). *Atlas Genet Cytogenet Oncol Haematol.*1998;2(2):60-61.

---